

Understanding the dynamics of fractional edge states with composite fermions

Dmitri B. Chklovskii and Bertrand I. Halperin

Physics Department, Harvard University, Cambridge, MA 02138

Fractional edge states can be viewed as integer edge states of composite fermions. We exploit this to discuss the conductance of the fractional quantized Hall states and the velocity of edge magnetoplasmons.

One of the most interesting developments in the physics of the quantized Hall effect was the introduction of the composite fermion approach.¹⁻⁴ It is realized through an exact transformation of the initial electron system to the system of composite fermions by attaching two flux quanta to each particle. The mean-field approximation to the new system already contains much of the physics related to electron-electron correlations. In particular it allows one to understand fractional quantized Hall states as integer quantized Hall states of composite fermions. Because of the presence of an excitation gap in the bulk, edge states play an important role in these systems. It is natural then to apply composite fermions to study fractional edge states. This has been done using the Hartree approximation for composite fermions^{5,6}, which yields ground state properties such as the density distribution and the number of edge channels for various filling factors. One has to go beyond the Hartree approximation in order to study the dynamics of the edge states, which involves excited states.

A significant progress in the understanding of edge states dynamics has been achieved by using the method of chiral bosonization⁷. However, this method relies on postulating the number and the direction of propagation of edge modes. The composite fermion approach allows one to derive this quantities microscopically. Once the velocities and interactions of edge modes are known they can be used as input parameters in the method of chiral bosonization.

In this paper we use the RPA to study fractional edge states. We begin by reviewing “gauge argument” and applying it to the system of composite fermions. This leads us to the introduction of the composite fermion electrochemical potentials which are very useful in calculating conductance of the system.

Following Laughlin⁸ and Halperin⁹ we consider an idealized clean 2DEG sample of the annular shape at filling factor ν . As the magnetic flux threading the hole is increased adiabatically by one flux quantum (ϕ_0) ν electrons are transferred from the outer to the inner edge of the sample. This leads to the total energy change given by $\nu(\xi_o - \xi_i)$, where $\xi_o - \xi_i$ is the difference between electrochemical potentials on the two edges. Since the work done on the system is given by $I\phi_0/c$, where I is the current around the annulus,

$$I = \nu(e/h)(\xi_o - \xi_i). \quad (1)$$

Now we rederive this result by using composite fermions. The electron system at filling factor $\nu = p/(2p + 1)$ is

viewed as the system of composite fermions at integer filling factor p . Naively, increasing the magnetic flux threading the annulus by one flux quantum will transfer p composite fermions between the outer and the inner edge. However the change in the flux $\Delta\phi$ enclosed by the annulus median comes from the flux carried by composite fermions as well as from the external source. Thus $\Delta\phi = (1 - 2x)\phi_0$, where x is the net number of composite fermions transferred between the edges. By relating the net transferred charge to the change in flux, using $x = p\Delta\phi/\phi_0$, we find that $x = p/(2p + 1)$ in agreement with the electron result.

It is helpful to introduce the composite fermion electrochemical potential, ξ^f . We define it as a change in the total energy resulting from adiabatically adding a composite fermion to the edge. The difference from the electron potential becomes clear in view of the previous discussion. The addition of a composite fermion to the edge involves turning on of the Chern-Simons flux, which adds or removes electrons from the edge. From the gauge argument we find the following relation between the two potentials:

$$\xi^f = \xi/(2p + 1) \quad (2)$$

This relation has been derived in a different way by Kirichenko and Johnson.¹⁰

If the edges of the annulus are sufficiently well separated currents associated with each edge can be defined. These currents must depend only on the electrochemical potential of the corresponding edge. Since the total current is given by Eq.1 the edge current is

$$I = \nu(e/h)\xi. \quad (3)$$

Expressing this result in terms of the composite fermion potential

$$I = p(e/h)\xi^f. \quad (4)$$

For the case $|p| > 1$, there will be several states at each edge, and in the ideal case one can define separate chemical potentials ξ^f on each edge state. We have assumed that there is a sufficient amount of scattering due to impurities or phonons, so that the states at a given edge are equilibrated, and there is a single chemical potential at each edge.

Now we consider an electromagnetic response of a single edge at a finite wavevector along the edge. The single edge approximation is justified if the wavelength is smaller than the distance between the edges. For a given

bulk filling factor there can be several magnetoplasmon modes at the edge if the impurity scattering is not too great. Here we restrict ourselves to the case of a sharp confining potential. In this case there is only one mode present for simple fractions $\nu = 1/(2k+1)$. We focus on the $\nu = 1$ case where the composite fermion result can be easily compared against the electron calculation. We can use previously derived formulas setting $p = -1$. Our goal is to find a pole in the response function and identify it with the magnetoplasmon mode.

We start by deriving the electron result. For frequencies smaller than the cyclotron frequency and wavelengths larger than the magnetic length eq.3 can be written in terms of Fourier harmonics

$$I(k, \omega) = (e/h)\xi(k, \omega). \quad (5)$$

The electrochemical potential can be represented as a sum of the external potential and the induced potential

$$\xi(k, \omega) = \xi^e(k, \omega) + \xi^i(k, \omega) \quad (6)$$

The induced electrochemical potential is proportional to the electron density at the edge

$$\xi^i(k, \omega) = n(k, \omega)\partial\xi/\partial n. \quad (7)$$

The coefficient $\partial\xi/\partial n$ is independent of k in the limit $k \rightarrow 0$, if the electron-electron interaction is short ranged; for unscreened Coulomb interactions we have $\partial\xi/\partial n \sim |\ln k|$.¹¹ For the wavelengths smaller than the length of the edge we invoke the continuity equation $n(k, \omega) = I(k, \omega)k/\omega$ to get

$$\xi^i(k, \omega) = I(k, \omega)(k/\omega)\partial\xi/\partial n \quad (8)$$

By substituting $I(k, \omega)$ from the eq.5 and solving for $\xi(k, \omega)$ we find

$$\xi(k, \omega) = \frac{\xi^e(k, \omega)}{1 - (k/\omega)(e/h)\partial\xi/\partial n} \quad (9)$$

Substituting this expression in eq.5 we obtain the full response function

$$I(k, \omega) = \frac{(e/h)\xi^e(k, \omega)}{1 - (k/\omega)(e/h)\partial\xi/\partial n} \quad (10)$$

At finite k and ω there is a pole in the response function, which can be identified with the magnetoplasmon mode. Because of the stability considerations $\partial\xi/\partial n$ has to be positive. Then the direction of propagation is in agreement with the classical magnetoplasmon result.¹¹

Attachment of two flux quanta transforms the $\nu = 1$ state into the $\nu = -1$ state with the effective magnetic field reversed. Therefore, on the mean field level one might expect that the composite fermion approach gives the reverse direction of propagation for the magnetoplasmon mode. In fact this is the result that comes out of

the Hartree approximation, where the induced Chern-Simons field is neglected. Let us demonstrate that including properly the Chern-Simons electric field gives the expected direction of propagation in the composite fermion approach. Eq.4 can be written in terms of Fourier harmonics

$$I(k, \omega) = -(e/h)\xi^f(k, \omega). \quad (11)$$

The composite fermion electrochemical potential ξ^f may be related to the electron potential ξ and the edge current I by

$$\xi(k, \omega) = \xi^f(k, \omega) + 2(h/e)I(k, \omega) \quad (12)$$

where the second term results from the Chern-Simons electric field produced by the current. [Eqs.11 and 12 imply Eq.2, as required] Substituting Eq. 12 in Eq. 6 we find

$$\xi^f(k, \omega) = \xi^e(k, \omega) - 2I(k, \omega)h/e + n(k, \omega)\partial\xi/\partial n \quad (13)$$

Using the continuity equation and expressing current in terms of the composite fermion electrochemical potential we get

$$\xi^f(k, \omega) = \xi^e(k, \omega) + 2\xi^f(k, \omega) - (k/\omega)\xi^f\partial\xi/\partial n \quad (14)$$

Solving this equation for ξ^f we find

$$\xi^f(k, \omega) = \frac{\xi^e(k, \omega)}{-1 + (k/\omega)(e/h)\partial\xi/\partial n} \quad (15)$$

Both of the terms in the denominator change signs here compared to Eq.9 thus giving the right direction of propagation for the magnetoplasmon mode. In Hartree approximation the Chern-Simons contribution to the electrochemical potential $2I(k, \omega)h/e$ is ignored thus yielding an incorrect direction of propagation.

For more complicated filling fractions ν , where there are two or more edge states at a single edge, it is necessary to take into account interactions between charge fluctuations associated with the different edge states. Similarly, if one considers wavelengths which are not large compared to the magnetic length and/or the geometric width of the edge, it is necessary to take into account more details of the electron wavefunctions. A natural approximation to use in this case is the RPA (or time dependent Hartree approximation) where the composite fermions are treated as non-interacting particles, driven by the space and time-dependent scalar and vector potentials. For a single edge state, the integral equation which results from the RPA reduces properly to the above equations in the limit where k and ω are small, giving the correct direction of propagation as expected.

We are grateful to A. Stern for helpful discussions. This work has been supported by the Harvard Society of Fellows and by NSF Grant No. DMR94-16910.

- ¹ J. K. Jain, Phys. Rev. Lett. **63**, 199 (1989).
- ² A. Lopez and E. Fradkin, Phys. Rev. B, **44**, 5246 (1991).
- ³ V. Kalmeyer and S.-C. Zhang, Phys. Rev. B **46**, 9889 (1992).
- ⁴ B. I. Halperin, P. A. Lee, and N. Read, Phys. Rev. B **47**, 7312 (1993).
- ⁵ L. Brey, Phys. Rev. B **50**, 11861 (1994).
- ⁶ D. B. Chklovskii, Phys. Rev. B **51**, 9895 (1995).
- ⁷ X.-G. Wen, J. Mod. Phys. B **6**, 1711 (1992).
- ⁸ R. B. Laughlin, Phys. Rev. B **23**, 5632 (1981).
- ⁹ B.I. Halperin, Phys. Rev. B **25**, 2185 (1982).
- ¹⁰ G. Kirczenow and B.L. Johnson, Phys. Rev. B **51**, 17579 (1995).
- ¹¹ V.A.Volkov, S.A. Mikhailov, Sov. Phys. JETP **67**, 1639 (1988).